

**Fisheries Oceanography of Yellowfin Tuna (*Thunnus
albacares*) in the Tasman Sea**



James Dell (BSc Hons.)

Submitted in partial fulfilment of the requirements for the degree of
Doctor of Philosophy in Quantitative Marine Science
University of Tasmania
November 2012

Cover illustration of yellowfin tuna, *Thunnus albacares*, by Roger Swainston © CSIRO
Marine Research, Australia

Abstract

Sustainable fishing is required to protect and maintain marine biodiversity and to ensure fisheries that are both economically viable and productive. Effective management of living marine resources requires well-informed decisions through an appreciation of past, present and future pressures. Understanding “why fish are caught where they are?” is the oldest question in fisheries research and is a central issue for the sustainable catch, management and conservation of marine resources. Here we look at the Australian longline fishery to better understand where yellowfin tuna (*Thunnus albacares*) are caught on the hooks set by the eastern tuna and billfish fishery (ETBF) in the Tasman Sea. We view the catch of this species in the context of the recent oceanography of the Tasman Sea, sixty years of fishermen’s knowledge and experience, a future climate scenario, declining stocks of other large pelagic predatory fish and an increasing demand in products derived from these species.

The physical environment directly influences the distribution, abundance, physiology and phenology of marine species. Relating species presence to physical ocean characteristics to determine habitat associations is fundamental to the management of marine species, however, direct observation of highly mobile animals in the open ocean, such as tunas and billfish, is challenging and expensive. As a result detailed data on habitat preferences using electronic tags has only been collected for the large iconic, valuable or endangered species. An alternative is to use commercial fishery catch data matched with historical ocean data to infer habitat associations. Using catch information from an Australian longline fishery and Bayesian hierarchical models we investigate the influence of environmental variables on the catch distribution of yellowfin tuna (Chapter 2). The focus was to understand the relative importance of space, time and ocean conditions on the catch of this pelagic predator. We found that pelagic regions with

elevated eddy kinetic energy, a shallow surface mixed layer and relatively high concentrations of chlorophyll *a* are all associated with high yellowfin tuna catch in the Tasman Sea. Time and space information, while important, were less informative than oceanic variables in explaining catch. An inspection of model prediction errors identified clumping of errors at margins of ocean features, such as eddies and frontal features, which indicate that these models could be improved by including representations of dynamic ocean processes which affect the catch of yellowfin tuna.

We use the same catch prediction model to consider where yellow fin tuna catches may occur in the context of a future climate scenario (Chapter 3). We used output from a global climate model (GCM) from the IPCC 2007 AR4 summit to produce predictions of surface ocean characteristics for the Tasman Sea in the 2060s. These data were used to initialize a biogeochemical model to create an ocean productivity product for the surface ocean that was equivalent to the chlorophyll *a* concentration as estimated by the ocean color SeaWiFS product. We use these products as inputs for the YFT catch prediction model to determine where YFT may be caught in the Tasman Sea in 2060s. We compare these predictions to those from the 1990s and the 2000s to show how the pattern of modelled YFT catches differ from those estimated by the model for the earlier time periods. Identifying possible shifts in the availability of YFT to commercial longlining over such long time period inform the construction of long-term goals upon which strategies for resource management; coastal infrastructure development and fleet management can be considered. This approach can also be applied at shorter time scales if biogeochemical downscaling is available.

Successful sustainable management of living marine resources can occur when the enhanced details of the resource, industry and market are thoughtfully integrated into the planning and implementation of management strategies. Engaging the fishing community

in the management process is a proven approach to the successful implementation of management strategies with sustainable outcomes. We report on a 2006 survey of the ETBF which recorded the perspectives of the resource users and cooperative managers regarding the location and catch of YFT in the Tasman Sea (Chapter 4). We show that the fishing community hold varied perspectives on the most influential ocean characteristics with respect to YFT catch and show how perspectives relate to the fishing region. Further work collecting, analysing and incorporating the opinions and knowledge of the fishermen of the ETBF into habitat and catch models is recommended as a direction for future work. Utilizing the qualitative information from fishers would minimise biases in the catch information, associated with the multispecies targeting and markets prices, and encourage better collaboration between fishermen, scientists and management for the sustainable future of the resource and fishing community of the east coast of Australia.

Overall, the work presented here show that YFT catches in the Tasman Sea can be partially explained by variation in the surface ocean environment. To achieve this goal, we used machine learning techniques to identify the most informative variables from the available ocean data and used a generalized linear model based on a hierarchical Bayesian framework to characterise the relationship between these variables and YFT catch. These techniques have not previously been used for this purpose in the Tasman Sea. The algorithms and model structures employed here provide a valid alternative to conventional habitat modelling techniques.

Declaration of Originality

This thesis contains no material which has been accepted for a degree or diploma by the University or any other institution, except by way of background information and duly acknowledged in the thesis, and to the best of my knowledge and belief no material previously published or written by another person except where due acknowledgement is made in the text of the thesis, nor does this thesis contain any material that infringes copyright.

Statement of Ethical Conduct

The research associated with this thesis abides by the international and Australian codes on the inclusion of human interactions in research. This research was conducted subject to the approval and guidelines of the Human Research Ethics Committee (Tasmania : reference H0008690).

James Dell 29 November 2012

Statement of publication and co-authorship

Publications produced as part of this thesis:

Dell, J., Wilcox, C.V. and Hobday, A.J. (2011) Estimation of yellowfin tuna (*Thunnus albacares*) habitat in waters adjacent to Australia's East Coast: making the most of commercial catch data. *Fisheries Oceanography*, **20**, 383-396.

Dell, J., Hobday, A.J., Chamberlain, M., Matear, R. and Wilcox, C.V. (in prep.) Potential impacts of climate change on the distribution of longline catches of yellowfin tuna (*Thunnus albacares*) in the Tasman Sea. *Climate Science*

Dell, J., Wilcox, C.V., Hobday, A.J. and Hindell M (in prep.) Fish Tales : can fishermen's expert knowledge improve models of tuna availability? (*Fish and Fisheries*)

Dell, J., Wilcox, C.V. and Hobday, A.J. (2007) *58th-Tuna-Conference-Proceedings*.
Lake Arrowhead, California

Dell, J., Wilcox, C.V. and Hobday, A.J. (2007) Aiding the habitat characterization of pelagic predators using expert opinion. *Annual meeting for the Society for Conservation Biology*. Port Elizabeth, Eastern Cape South Africa

The following people and institutions contributed to the publication of the work undertaken as part of this thesis:

- *Chris Wilcox (CSIRO Marine and Atmospheric Research), Alistair Hobday (CSIRO Marine and Atmospheric Research) and Mark Hindell (Institute of Marine and Antarctic Studies, University of Tasmania) assisted with guidance and supervision in all aspects of the PhD and producing publishable quality manuscripts*
- *Richard Matear and Matt Chamberlain (both of CSIRO Marine and Atmospheric Research) provided assistance with the usage of data from dynamically downscaled Global Circulation Models*
- *Robert Campbell (CSIRO Marine and Atmospheric Research) provided assistance with the usage of commercial fisheries logbook information and guidance with the management and suitable applications of these data*
- *Campbell Davies (CSIRO Marine and Atmospheric Research) provided guidance on fisheries stakeholder interactions and resource management aspects of the research and in producing publishable quality manuscripts.*

We the undersigned agree with the above stated “proportion of work undertaken” for each of the above published (or in preparation) peer-reviewed manuscripts contributing to this thesis.

James Dell (Candidate)

Mark Hindell (Candidate’s Supervisor)

Tom Trull (Head of school)

Acknowledgements

First of all, I must thank my supervisors Chris Wilcox, Alistair Hobday and Mark Hindell, whom I am deeply grateful for their expertise, broad vision, insight and patience. In particular, thanks to Alistair for encouraging me to undertake this challenging project. Without his successful guidance and encouragement through my honors year, I would have never considered this path. His experience, extraordinary productivity and breath of engagement in the marine realm were an inspiration. Thanks to Chris for his unique vision and excitement for new approaches in understanding ecological processes, particularly when confronted with challenging data sets. Special thanks must be extended for his efforts and patience engaged when introducing (and re-introducing) me to realm of Bayesian statistics and the power of machine learning. Finally, thanks to Mark for his sincere and cheerful dedication to and belief in his students. His academic experience, positivity, incisive guidance and can-do-attitude were of great aid and comfort at times when the going was particularly tough. I was well supported by both the joint UTas/CSIRO Quantitative Marine Science program and CMAR's Wealth from Oceans Flagship. These initiatives are helping to foster the future great achievements in marine science in Australia and beyond.

Thanks also to Richard Coleman, Tom Trull and Simon Wotherspoon for their perspectives, guidance and wisdom in their roles as QMS co-ordinators. Campbell Davies played many roles in this saga, from mentor and motivator to humorist, surfing conspirator and close friend. I thank him specifically for his continued interest and belief in my project, particularly at the times when I had lost mine. His perspectives, positivity and wisdom were key to me continuing to push through towards the completion of this work and this thesis. I also benefitted from the tolerance and patience of Toby Patterson when seeking his knowledge on the nuances of R programming, and his perspectives on

statistics and rigorous scientific method. John Gunn provided five sage-like words early on, which I have continually reflected on and, in the end found to be true. Robert Campbell provided great insight and guidance in relation to the dealing with the ETBF logbook information provided by fishermen via AFMA. He also reviewed an early iteration of the first data chapter and helped in guiding it towards the published form. Beyond their kind encouragement and compassion in the corridor and at the kettle, Karen Evans and Marinelle Basson provided incredibly thorough reviews of chapters and very helpful perspectives on the ecology of pelagic predators. Their critical input was instrumental in improving the rigor and clarity of some of the following chapters. I must also thank the ex-corporeal “Pelagics group” at CSIRO Marine and Atmospheric Research (CMAR) for being one of the most conscientious, humble, inspiring, fiercely intelligent and sincerely compassionate bunch of people I have ever had the pleasure to meet. Based purely on my experience within this group of people, furthering a career in science is an attractive option – even if only to associate with others whom share these same qualities. You are all amazing and surely some of the best global citizens anyone is likely to meet.

I am grateful to Mark Mangel and Don “Diego” Croll, who welcomed me to their labs, provided desks, imparted some wise words and perspectives and were just-all-round-inspirational while I visited UCSC briefly in 2006. I extend heartfelt thanks to Stacey Buckelew for her friendship, inspiration, belief and tangential influences during my time as a graduate student. Straight lines are boring. Life and research should be a winding path. I finally JFDI and will be sure to remember your safety advice on all future expeditions.

A warm thanks to the fellow inhabitants of the student dens at the Institute of Marine and Antarctic Studies (UTAS) and CMAR, for sharing the joys and frustrations that only a PhD can evoke. Thanks also for the shared cakes and coffee runs that helped to break up the day and reminded me that I was once a social creature.

Special thanks are extended to the skippers of the ETBF who graciously volunteered their time and experience in answering questions that sought to dig out the invaluable information they have accumulated from many days and nights at sea.

I also wish to thank all those close and dear to me for their support and love during perhaps the hardest challenge I have faced. Mum thanks for the love and good vibes always. Dad, thanks for the inherited stubbornness and dedication. I am so very happy to share this achieve with you both.

And to the amazing Maia! For all the love, laughter and tolerance (massage, physio, acupuncture, culinary genius, domestic bliss and for finding the shack!). All your gorgeous grace, unending generosity, enduring patience and deep love kept me on track in the final stretch. Now we can finally go on holiday!

Table of Contents

Abstract	i
Declaration of Originality	iv
Statement of Ethical Conduct.....	iv
Statement of publication and co-authorship	v
Acknowledgements	vii
Table of Contents	x
List of Figures	xii
List of Tables.....	xv
CHAPTER 1.....	1
General Introduction	1
The Global Context for tuna and tuna fisheries	1
The Tasman Sea and Eastern Australian Tuna and Billfish Fishery.....	6
Spatial dynamics of YFT in the Tasman Sea	12
CHAPTER 2.....	18
Abstract	19
Introduction	20
Methods.....	24
Fisheries Data.....	24
Ocean data	24
Variable Importance using Random Forest Algorithm	26
Hierarchical Modeling using Bayesian mixture models	26
Estimation of model variance and errors.....	27
Model structure and selection	28
Residual analysis	30
Results	31
Distribution of longline fishing effort	31
Preliminary variable selection.....	31
Model selection	35
Discussion	41
CHAPTER 3.....	48
Introduction	49
Models, data and Methods	52
Catch and effort data.	52
Modelling approaches	52
Approach 1 : Predicted future YFT catch based on observations from recent past ...	55
Approach 2 : Predicted change in YFT catch from modelled climate scenarios	57
Results	59
Approach 1 : Hot and cold seasons from recent historical data: 2000 to 2010.....	59
YFT catch rate predictions for recent climate.	60
Reviewing spatial ocean data from “simulated 2000s” and “simulated 2060s”	
npzdsGCM experiments.....	64
Approach 2 : Differences in YFT catch rates between the simulated 2000s and the	
simulated 2060s.....	66
Discussion	70
CHAPTER 4.....	78
Fish Tales: can fisherman’s expert knowledge improve models of tuna availability? ..	78
Introduction	79

Methodology	80
Data collection	81
Collation of elicited information on ocean characteristics	82
Multivariate Primer analyses.	83
Assessing associations between fishers' interview responses and characteristics of their fishing strategy and success.....	85
Results.....	86
Interview Responses	86
Multivariate analyses of interview responses	87
Fishing locations and catch efficiency of interview cluster groups	90
Multivariate analysis of fishermen's catch performance data	92
Fishing locations of cluster groups based on catch performance metrics.....	93
Multivariate analysis of fishermen's primary fishing zone	93
Maps of skippers fishing zone groupings	96
Maps of skipper grouped based on the region of their homeport	97
Associations between fishers' interview responses and characteristics of their fishing strategy and success	100
Discussion	103
Why might Fishermen show a diverse range of perceptions on the specifics of YFT habitat?.....	104
CHAPTER 5	110
Synthesis, future directions and conclusions	110
Spatial models of large pelagic predators	113
Potential impacts of climate extremes on future distributions.....	115
Incorporating expert knowledge	119
Future focus and directions	122
Conclusions.....	125
REFERENCES	126
Appendix –.....	143
Outline of the interview	143
Compilation of selected interview responses.....	156
Statistical details and results	158